

# SMALLPOX VACCINATION PROGRAM

THE DANGER

THE VACCINE

THE PRIORITY

THE STRATEGY

adverse event info

education toolkit

resource center

my vaccination

smallpox overview



search

sitemap

## ADVERSE EVENT

- [Safety Panel F](#)
- [Cardiac Disease](#)
- [Types of Reactions](#)
- [What do I do?](#)
- [How do I learn?](#)
- [Pregnancy Considerations](#)
- [Overview](#)
- [VAERS Form](#)
- [Safety Summary](#)

## SMALLPOX VACCINATION SAFETY SUMMARY

### DoD Smallpox Vaccination Program

Safety Summary, May 02, 2005

#### Background:

On December 13, 2002, the President directed smallpox vaccinations for selected military personnel, government workers, and contracted workers. DoD vaccinations began immediately for emergency response personnel and hospital staff members. Comprehensive training programs in vaccination technique, infection-control safeguards, screening and education methods, adverse event monitoring, product storage and handling, aggressively launched in October 2002, made immediate vaccination possible. In early January 2003, DoD began smallpox vaccinations of selected US military forces, a emergency-essential civilians and contractors deployed or deploying in support of U.S. Central Command missions.

#### Program Status:

DoD operational forces and healthcare workers vaccinated against smallpox: over 800,000. Detail of the DoD program appeared in the June 25, 2003 issue of the Journal of the American Medical Association (JAMA). The abstract of that article appears after the following summation of the program's current status.

In the 28.5 months between 13 Dec 02 and 02 May 05, the DoD administered more than 800,000 smallpox vaccinations. Most adverse events occurred at rates below historical rates.

- 97 cases of myo-pericarditis developed after smallpox vaccination. These cases have been followed carefully to evaluate their recovery, at over 30 hospitals in 25 States and several countries overseas. Detailed follow-up cardiac testing is available in 64 cases: all 64 had normal electrocardiograms (EKGs), and normal treadmill stress test results, suggesting a high rate of recovery. [See Journal of the American College of Cardiology. 2004 (Jul 7); 44(1):201-5.]
- Another 16 cases of "ischemic" heart disease (such as heart attacks, atherosclerosis, or angina) occurred within 6 weeks after smallpox vaccination. This number of cases is similar to what normally occurs among unvaccinated military personnel of similar age.
- In no case was vaccinia (the live virus in the vaccine) transmitted from one person to another in the work place. Among 27,700 smallpox-vaccinated health-care workers, there were no cases of transmission of vaccinia from worker to patient.
- 53 cases (30 lab-confirmed) of contact transfer of vaccinia virus have occurred, principally to spouses and adult intimate contacts. During education, we repeat the warning "Don't let your guard down at home."
- No cases of eczema vaccinatum or progressive vaccinia occurred. This indicates our education and exemption process is working well.
- The total number of treatments with vaccinia immune globulin (VIG) remains at three: one burn and two ocular patients.
- 43 cases of possible generalized vaccinia were treated, primarily as outpatients. Few of those cases meet a strict case definition.
- Seven deaths due to disease after vaccination have been reviewed: one following an acute lupus illness may have been caused by vaccination, based on review by two independent panels of civilian physicians. Additional information on this case appears at <http://www.smallpox.mil/event/panelreport.asp>; other deaths involved the following diagnoses (one each, except as noted): myocardial infarction, atherosclerotic coronary vascular disease (ASCVD, two), pulmonary embolism, heat injury, and benzodiazepine overdose. These deaths were judged unrelated to vaccination, based on individual circumstances such as preexisting disease, incidence among unvaccinated people, and lack of physical evidence of

implicate a vaccine.

NOTE: All appropriate program information is provided regularly to federal health authorities, including safety-surveillance data.

The June 25, 2003, issue of the Journal of the American Medical Association (JAMA) summarizes the experience with smallpox vaccination to date. Below appears the abstract of this article:

**US Military Smallpox Vaccination Program Experience** John D. Grabenstein, RPh, PhD  
William Winkenwerder, Jr., MD, MBA

**Context.** The United States recently implemented smallpox vaccination of selected military personnel as part of a national program of preparedness against use of smallpox as a biological weapon. The return of smallpox vaccinations raises important questions regarding implementation and safety.

**Objective.** To describe the US military smallpox vaccination program.

**Design.** Descriptive study of the vaccination program from its inception on December 13, 2002, to May 28, 2003.

**Setting.** US Department of Defense (DoD) fixed and field medical treatment facilities on multiple continents and ships at sea.

**Subjects.** US service members and DoD civilian workers eligible for smallpox vaccination.

**Main Outcome Measures.** Numbers of vaccinations and rates of vaccination exemptions, symptomatic adverse events. Data was collected via reports to headquarters and rigorous surveillance for sentinel events.

**Results.** In 5.5 months, the DoD administered 450,293 smallpox vaccinations (70.5% primary vaccinees and 29.5% revaccinees). In 2 settings, 0.5% and 3.0% of vaccine recipients needed short-term sick leave. Most adverse events occurred at rates below historical rates. One case of encephalitis and 3 cases of acute myopericarditis developed after vaccination; all cases recovered. Among 19,461 workers with no clinical contact, there were no cases of transmission of vaccinia from worker to patient, no case of eczema vaccinatum or progressive vaccinia, and no attributed deaths.

**Conclusions.** Mass smallpox vaccinations can be conducted safely with very low rates of serious adverse events. Program implementation emphasized human factors: careful staff training, contraindication screening, recipient education, and attention to bandaging. Our experience suggests broad smallpox vaccination programs may be implemented with fewer serious adverse events than previously believed. JAMA. 2003;289:3278-3282 [www.jama.com](http://www.jama.com)

Full article at: <http://jama.ama-assn.org/cgi/reprint/289/24/3278.pdf>

For more discussion of the myo-pericarditis cases, see: <http://jama.ama-assn.org/cgi/reprint/289/24/3278.pdf>

For the accompanying editorial, see: <http://jama.ama-assn.org/cgi/reprint/289/24/3306.pdf>

Perspective:

"Within DoD, we know a lot now about smallpox vaccine. We now have current, good experience with using it. The remarkable news about this program is that our adverse events do not reflect what has been told us to expect. With proper care and attention, smallpox vaccine can be administered to large numbers of people safely. Those large numbers of vaccinated citizens discourage an enemy from using smallpox as a weapon."

- William Winkenwerder, Jr., MD, Assistant Secretary of Defense for Health Affairs

[PDF version \(March 01,2005\)](#)

[PDF version \(October 14,2004\)](#)

[PDF version \(August 25,2004\)](#)

[PDF version \(June 16,2004\)](#)

[PDF version \(May 14,2004\)](#)

[PDF version \(March 04, 2004\)](#)

[PDF version \(February 11, 2004\)](#)

[PDF version \(December 12, 2003\)](#)

[PDF version \(November 19, 2003\)](#)

[PDF version \(October 22, 2003\)](#)

[PDF version \(June 25, 2003\)](#)

[PDF version \(May 16, 2003\)](#)

[PDF version \(March 31, 2003\)](#)

[PDF version \(February 12, 2003\)](#)

[PDF version \(January 31, 2003\)](#)

Last Updated:  
15 Jul 05

